


# ANNALES MATHÉMATIQUES



## BLAISE PASCAL

SALAH HAJJI

Stochastic fractional partial differential equations driven  
by Poisson white noise

Volume 15, n° 1 (2008), p. 43-55.

[http://ambp.cedram.org/item?id=AMBP\\_2008\\_\\_15\\_1\\_43\\_0](http://ambp.cedram.org/item?id=AMBP_2008__15_1_43_0)

© Annales mathématiques Blaise Pascal, 2008, tous droits réservés.

L'accès aux articles de la revue « Annales mathématiques Blaise Pascal » (<http://ambp.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://ambp.cedram.org/legal/>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

*Publication éditée par le laboratoire de mathématiques  
de l'université Blaise-Pascal, UMR 6620 du CNRS  
Clermont-Ferrand — France*

**cedram**

Article mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques  
<http://www.cedram.org/>

# Stochastic fractional partial differential equations driven by Poisson white noise

SALAH HAJJI

## Abstract

We study a stochastic fractional partial differential equations of order  $\alpha > 1$  driven by a compensated Poisson measure. We prove existence and uniqueness of the solution and we study the regularity of its trajectories.

*Équations aux dérivées partielles fractionnaires stochastiques dirigées par un bruit poissonnien*

## Résumé

On étudie une équation aux dérivées partielles stochastiques fractionnaires d'ordre  $\alpha > 1$  dirigée par une mesure de Poisson compensée. On montre l'existence et l'unicité de la solution et on étudie la régularité de ses trajectoires.

## 1. Introduction

In recent years fractional calculus knew a great progress considering their applications in different fields of science, including numerical analysis, physics, engineering, biology, economics and finance.

Partial differential equations is one of the mathematical domains where the fractional calculus is strongly used (see Podlubny [8]). However, few publications treat stochastic partial differential equations involving fractional derivatives. Most of them investigate evolution type equations, driven by a fractional power of the Laplacian (see [3], [10]). These operators generate symmetric stable semigroups when the order of derivation is less than 2. Our idea is inspired by the paper of Debbi and Dozzi [5] in which the authors studied a nonlinear stochastic fractional partial differential equations of high order containing also derivatives of entire order and perturbed by

---

*Keywords:* Stochastic partial differential equations; fractional derivative operator; Poisson measure.

*Math. classification:* 26A33, 60H15.

Gaussian white noise.

The aim of this paper is to generalize the result of [1] to nonlinear stochastic fractional partial differential equations. We study the existence, uniqueness and regularity of the solution of the following one-dimensional stochastic fractional partial differential equation, formally given by

$$\begin{aligned} \frac{\partial u}{\partial t}(t, x, \omega) &= D_\alpha^\delta u(t, x, \omega) + f(t, u(t, x)) + \int_{\mathcal{U}} g(t, u(t, x), z) \tilde{N}(dt, dx, dz) \\ u(0, x) &= u_0(x) \end{aligned} \quad (1.1)$$

for  $t \in (0, +\infty)$ ,  $x \in \mathbb{R}$ , where  $(\mathcal{U}, \mathcal{B}(\mathcal{U}), q)$  is a  $\sigma$  finite measure space,  $D_\delta^\alpha$  is the fractional differential operator with respect to the spacial variable,  $f : (0, +\infty) \times \mathbb{R} \longrightarrow \mathbb{R}$  and  $g : (0, +\infty) \times \mathbb{R} \times \mathcal{U} \longrightarrow \mathbb{R}$  are measurable.

One rigorous formulations of the equation (1.1) can be given by the following integral equation

$$\begin{aligned} u(t, x) &= \int_{\mathbb{R}} G_\alpha(t, x - y) u_0(y) dy \\ &+ \int_0^t \int_{\mathbb{R}} G_\alpha(t - s, x - y) f(s, u(s, y)) dy ds \\ &+ \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t - s, x - y) g(s, u(s, y), z) \tilde{N}(ds, dy, dz) \end{aligned} \quad (1.2)$$

for  $t \in [0, +\infty)$  and  $x \in \mathbb{R}$ , where  $\{G_\alpha(t, x), x \in \mathbb{R}, t \in [0, +\infty)\}$  stands for the fundamental solution of the operator  $D_\delta^\alpha$ .

Some equations similar to equation (1.1) are studied also in [6] and [2], but with the Laplacian operator in place of the operator  $D_\delta^\alpha$ .

We assume that the functions  $f$  and  $g$  satisfy the following growth and Lipschitz conditions:

(H)

For all  $T > 0$ , there exists a constant  $K = K_T$ , such that for all  $0 \leq t \leq T$ ,  $x, y \in \mathbb{R}$ ,  $z \in \mathcal{U}$

- $|f(t, x)|^2 + \int_{\mathcal{U}} |g(t, x, z)|^2 q(dz) \leq K_T(1 + |x|^2)$
- $|f(t, x) - f(t, y)|^2 + \int_{\mathcal{U}} |g(t, x, z) - g(t, y, z)|^2 q(dz) \leq K_T|x - y|^2$ .

The remaining of the paper is structured as follows. In Section 2, we give the definitions of stochastic integral with respect to  $\tilde{N}(ds, dy, dz)$  and the derivative operator  $D_\delta^\alpha$ , then we recall some inequalities satisfied by the fundamental solution  $G_\alpha$  and we precise the notion of solutions

of the equation (1.1). Section 3 is dedicated to the proof of existence and uniqueness of the solution. In Section 4 we prove the spatial Hölder continuity of the solution.

## 2. Preliminaries and definitions

Let  $(\Omega, \mathcal{F}, P)$  be a probability space equipped with a filtration  $\{\mathcal{F}_t\}$  satisfying the usual conditions. Let  $(\mathcal{U}, \mathcal{B}(\mathcal{U}), q)$  be a  $\sigma$  finite measure space and let  $N$  be a Poisson measure on  $\mathbb{R}_+ \times \mathbb{R} \times \mathcal{U}$  with intensity measure  $\nu(ds, dx, dz) = ds dx q(dz)$ . The compensated Poisson measure is denoted by  $\tilde{N} = N - \nu$ .

Now, we define the stochastic integrals we will use.

### 2.1. Stochastic integral with respect to $\tilde{N}(ds, dy, dz)$

Let us introduce the following class:

$$F_{pred} := \left\{ h(t, y, z, \omega) : h \text{ is } \mathcal{F}_t \text{ predictable and } \forall t > 0, \right. \\ \left. E \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} |h(s, y, z)|^2 \nu(ds, dy, dz) < \infty \right\}$$

it is known (see Ikeda and Watanabe [7]) that for any  $t > 0$ , the stochastic integral  $\int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} h(s, y, z) \tilde{N}(ds, dy, dz)$ , for  $h \in F_{pred}$ , can be well defined. The stochastic integral has the following isometry property

$$E \left\{ \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} h(s, y, z) \tilde{N}(ds, dy, dz) \right\}^2 = \\ E \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} |h(s, y, z)|^2 \nu(ds, dy, dz).$$

Thus for any  $t > 0$ ,  $\int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} h(s, y, z) \tilde{N}(ds, dy, dz) \in L^2(\Omega)$ .

This integral can be extended to a more general class  $F$  of integrands without the predictable property, in the following manner :

A function  $h$  is said to be of class  $F$  if is  $\mathcal{F}_t$  adapted, and there exists a sequence  $h_n \in F_{pred}$  such that for any  $t > 0$

$$E \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} |h_n(s, y, z) - h(s, y, z)|^2 \nu(ds, dy, dz) \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

For  $h \in F$ , for any  $t > 0$ ,  $\int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} h(s, y, z) \tilde{N}(ds, dy, dz)$  is defined as the  $L^2(\Omega)$ -limit of the following Cauchy sequence

$$\left\{ \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} h_n(s, y, z) \tilde{N}(ds, dy, dz) \right\}_{n \in \mathbb{N}}$$

(see Albeverio *et al.* [1] for more details).

## 2.2. Definition of the operator $D_\delta^\alpha$

The fractional differential operator  $D_\delta^\alpha$  is an extension of the inverse of the generalized Riesz-Feller Potential when  $\alpha > 2$ . It is given for  $\alpha > 0$  by

**Definition 2.1.** The fractional differential  $D_\delta^\alpha$  is given by

$$D_\delta^\alpha \varphi = \mathcal{F}^{-1} \{ \psi_\alpha \cdot \mathcal{F} \varphi \}$$

where

$$\psi_\alpha(\lambda) = -|\lambda|^\alpha e^{-i\delta \frac{\pi}{2} \operatorname{sgn}(\lambda)}.$$

$|\delta| \leq \min\{\alpha - [\alpha]_2, 2 + [\alpha]_2 - \alpha\}$ ,  $[\alpha]_2$  is the largest even integer less or equal to  $\alpha$ , and  $\delta = 0$  when  $\alpha \in 2\mathbb{N} + 1$ .  $\mathcal{F}$  (respectively  $\mathcal{F}^{-1}$ ) is the Fourier (respectively Fourier inverse) transform.

The operator  $D_\delta^\alpha$  is a closed, densely defined operator on  $L^2(\mathbb{R})$  and it is the infinitesimal generator of a semigroup which is in general not symmetric and not a contraction. It is selfadjoint only when  $\delta = 0$  and in this case, it coincides with the fractional power of the Laplacian. When  $\alpha = 2$  it is the Laplacian itself (see [4, 5]).

The Green function  $G_\alpha(t, x)$  associated to the equation (1.1) is the fundamental solution of the Cauchy problem

$$\begin{cases} \frac{\partial}{\partial t} G(t, x) = D_\alpha^\delta G(t, x), & t > 0, x \in \mathbb{R}, \\ G(0, x) = \delta_0(x). \end{cases}$$

Using Fourier's calculus we obtain

$$G_\alpha(t, x) = \mathcal{F}^{-1} \{ e^{\delta \psi_\alpha(\lambda)t}; x \} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \exp(-i\lambda x - t|\lambda|^\alpha e^{-i\delta \frac{\pi}{2} \operatorname{sgn}(\lambda)}) d\lambda.$$

Let us recall some well-known properties, (see [5]), of the Green kernel  $G_\alpha$  which will be used later.

**Lemma 2.2.** For  $\alpha \in (0, +\infty)$

(1)  $\int_{\mathbb{R}} G_{\alpha}(t, x) dx = 1.$

(2)  $G_{\alpha}(t, x)$  satisfy the semigroup property, i.e. for  $0 < s < t$

$$G_{\alpha}(t + s, x) = \int_{\mathbb{R}} G_{\alpha}(t, y) G_{\alpha}(s, x - y) dy.$$

(3) For  $0 < \alpha \leq 2$ , the function  $G_{\alpha}(t, \cdot)$  is the density of a Lévy stable process in time  $t$ .

(4) For fixed  $t$ ,  $G_{\alpha}(t, \cdot) \in S^{\infty} = \{f \in C^{\infty} \text{ and } \frac{\partial^k}{\partial x^k} f \text{ is bounded and tends to zero when } |x| \text{ tend to } \infty, \forall k \in \mathbb{N}\}.$

(5)  $\frac{\partial^n G_{\alpha}}{\partial x^n}(t, x) = t^{-\frac{n+1}{\alpha}} \frac{\partial^n G_{\alpha}}{\partial \xi^n}(1, \xi) \Big|_{\xi=t-\frac{1}{\alpha}x},$  for all  $n \geq 0$ .

(6) Let  $\alpha \in (1, +\infty)$ . Then there exists a constant  $K_{\alpha}$  such that

- $|G_{\alpha}(1, x)| \leq K_{\alpha}(1 + |x|^{1+\alpha})^{-1},$
- $|G_{\alpha}^n(1, x)| \leq K_{\alpha} \frac{1+|x|^{\alpha+n-1}}{(1+|x|^{\alpha+n})^2}.$

(7) For  $\alpha \in (1, +\infty)$ , for any fixed  $n \in \mathbb{N}$ , and  $T \geq 0$ , for  $\gamma$  such that  $\frac{1}{\alpha+n+1} < \gamma < \frac{\alpha+1}{n+1},$

$$\int_0^T \int_{-\infty}^{+\infty} \left| \frac{\partial^n G_{\alpha}}{\partial y^n}(s, y) \right|^{\gamma} dy ds < \infty.$$

In this paper, we restrict ourselves to the case  $\alpha > 1$  in order that  $\int_0^T \int_{\mathbb{R}} G_{\alpha}^2(t, x) dx ds < \infty$  (see Lemma 2.2, property (7)).

Let us now give a precise formulation of solutions for Equation (1.1).

**Definition 2.3.** Let  $0 < T < \infty$ . A measurable stochastic field

$$\{u(t, x), t \in [0, T], x \in \mathbb{R}\}$$

is said to be a mild solution of the equation (1.1) on  $[0, T]$  if

- $u(t, x)$  is adapted for every  $x$ .
- $\{u(t, x)\}$ , as a family of  $L^2(\Omega, \mathcal{F}, P)$ -valued random variables, is right continuous and has left limits in the variable  $t \in [0, T]$ , namely,

$$u(t-, x, \cdot) = L^2(\Omega) - \lim_{s \uparrow t} u(s, x, \cdot), \quad t \in [0, T].$$

In this paper we simply call such a  $u$  *modified càdlàg* in  $t$  (after the French acronym).

- Equation (1.2) holds a.s. for all  $t \in [0, T]$  and  $x \in \mathbb{R}$ .

A solution  $u$  on  $[0, T]$  is said to be  $L^p(\Omega)$ - bounded if

$$\sup_{t \leq T} \sup_{x \in \mathbb{R}} E|u(t, x)|^p < \infty.$$

All positive constants appearing in this paper are called  $C$ . They may change from one line to the next one.

### 3. Existence and uniqueness

The main result of this section is the following

**Theorem 3.1.** *Let  $p \geq 2$  and  $T > 0$  be fixed. Under condition (H) and the assumption that the initial condition  $u_0$  is  $L^p(\Omega)$  bounded, equation (1.1) has a unique mild solution  $u$  on  $[0, T]$  which is  $L^p(\Omega)$  bounded.*

For the proof we need the following lemma.

Fix  $T > 0$  and let

$$\mathcal{E} := \left\{ \{v(t, x), t \in [0, T], x \in \mathbb{R}\} : v \text{ is measurable, } \mathcal{F}_t \text{ adapted,} \right. \\ \left. \text{modified càdlàg in } t \text{ such that, } \sup_{t \leq T} \sup_{x \in \mathbb{R}} E|v(t, x)|^p < \infty \right\}.$$

**Lemma 3.2.** *Let  $v \in \mathcal{E}$  and set*

$$X(t, x) := \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t-s, x-y) g(s, v(s-, y), z) \tilde{N}(ds, dy, dz),$$

$$Y(t, x) := \int_0^t \int_{\mathbb{R}} G_\alpha(t-s, x-y) f(s, v(s, y)) dy ds.$$

*Then  $X$  and  $Y$  belong to  $\mathcal{E}$ .*

*Proof.* Let us first show that  $X$  is well defined.

For any fixed  $t \in [0, T]$ , let, for  $(s, y, z) \in [0, t] \times \mathbb{R} \times \mathcal{U}$ ,

$$h_{t,x}(s, y, z) := G_\alpha(t-s, x-y) g(s, v(s-, y), z),$$

$$v^n(s, y) := v(0, y) 1_{\{0\}}(s) + \sum_{k=0}^{2^n-1} v\left(\frac{kt}{2^n}, y\right) 1_{\left(\frac{kt}{2^n}, \frac{(k+1)t}{2^n}\right]}(s), \quad n \in \mathbb{N},$$

and

$$h_{t,x}^n(s, y, z) := G_\alpha(t - s, x - y)g(s, v^n(s, y), z).$$

Clearly  $v^n$  is  $\mathcal{F}_s$  predictable, so  $h^n$  is also  $\mathcal{F}_s$  predictable. Let us show that  $h^n \in F_{pred}$ .

By condition(H) we have :

$$\begin{aligned} & E \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} |h_{t,x}^n(s, y, z)|^2 \nu(ds, dy, dz) \\ & \leq K_T \int_0^t \int_{\mathbb{R}} G_\alpha(t - s, x - y)^2 (E|v^n(s, y)|^2 + 1) dy ds < \infty. \end{aligned}$$

Thus  $h_{t,x}^n \in F_{pred}$ . On the other hand, by the condition (H) we have

$$\begin{aligned} & E \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} |h_{t,x}^n(s, y, z) - h_{t,x}(s, y, z)|^2 \nu(ds, dy, dz) \\ & \leq K_T \int_0^t \int_{\mathbb{R}} G_\alpha(t - s, x - y)^2 E|v^n(s, y) - v(s-, y)|^2 ds dy \end{aligned}$$

By the definition of  $v^n$  and the fact that  $v$  is modified càdlàg we have

$$E|v^n(s, y) - v(s-, y)|^2 \xrightarrow{n \rightarrow \infty} 0, \quad \text{for all } (s, y) \in [0, t] \times \mathbb{R}$$

and

$$E|v^n(s, y) - v(s-, y)|^2 \leq 4 \sup_{r \leq T} \sup_{x \in \mathbb{R}} E|v(r, x)|^2, \quad \forall (s, y, n) \in [0, t] \times \mathbb{R} \times \mathbb{N}.$$

Furthermore by Lemma 2.2, property (7), we have

$$\int_0^t \int_{\mathbb{R}} G_\alpha(t - s, x - y)^2 ds dy < \infty.$$

Then by the Lebesgue's dominated convergence theorem we obtain that

$$\int_0^t \int_{\mathbb{R}} G_\alpha(t - s, x - y)^2 E|v^n(s, y) - v(s-, y)|^2 ds dy \xrightarrow{n \rightarrow \infty} 0.$$

Hence  $h_{t,x} \in F$  for any fixed  $t \in [0, T]$ , so the integral

$\int_0^{t+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t - s, x - y)g(s, v(s-, y), z)\tilde{N}(ds, dy, dz)$  is well defined.

Using the same techniques, as in [1], we show that  $X$  is  $\mathcal{F}_t$  adapted and modified càdlàg.



S. HAJJI

Further more, by Burkholder's inequality and condition (H), we have

$$\begin{aligned}
E|X(t, x)|^p &\leq CE \left( \int_0^t \int_{\mathbb{R}} \int_{\mathcal{U}} G_{\alpha}^2(t-s, x-y) g^2(s, v(s-, y), z) ds dy q(dz) \right)^{\frac{p}{2}} \\
&\leq CE \left( \int_0^t \int_{\mathbb{R}} G_{\alpha}^2(t-s, x-y) (1 + v^2(s-, y)) ds dy \right)^{\frac{p}{2}} \\
&\leq C(1 + \sup_{s \leq T} \sup_{x \in \mathbb{R}} E|v(s, x)|^p) \int_0^T \int_{\mathbb{R}} G_{\alpha}^2(t-s, x-y) ds dy)^{\frac{p}{2}}
\end{aligned}$$

By Lemma 2.2, property (7), the last integral is finite. Then

$$\sup_{t \leq T} \sup_{x \in \mathbb{R}} E|X(t, x)|^p < \infty$$

Finally  $X \in \mathcal{E}$ .

It is well known that  $Y$  is measurable  $\mathcal{F}_t$  adapted and modified càdlàg. By Hölder inequality with respect to the measure  $|G_{\alpha}(t-s, x-y)| ds dy$  and the growth assumption on  $f$ , we have

$$\begin{aligned}
E|Y(t, x)|^p &\leq E \left( \int_0^t \int_{\mathbb{R}} |G_{\alpha}(t-s, x-y)| f^p(s, v(s, y)) ds dy \right) \\
&\quad \times \left\{ \int_0^t \int_{\mathbb{R}} |G_{\alpha}(t-s, x-y)| ds dy \right\}^{p-1} \\
&\leq CE \int_0^t \int_{\mathbb{R}} |G_{\alpha}(t-s, x-y)| (1 + v^p(s, y)) ds dy \\
&\leq C(1 + \sup_{s \leq T} \sup_{x \in \mathbb{R}} E|v(s, x)|^p) \int_0^T \int_{\mathbb{R}} |G_{\alpha}(t-s, x-y)| ds dy.
\end{aligned}$$

So

$$\sup_{t \leq T} \sup_{x \in \mathbb{R}} E|Y(t, x)|^p < \infty.$$

and finally  $Y \in \mathcal{E}$ . □

*Proof of Theorem 3.1.*

• Proof of the existence: Let the sequence  $(u_n, n \geq 1)$  be given by:

$$u_1(t, x) = \int_{\mathbb{R}} G_{\alpha}(t, x-y) u_0(y) dy$$

FRACTIONAL SPDES DRIVEN BY POISSON WHITE NOISE

$$\begin{aligned} u_{n+1}(t, x) &= u_1(t, x) + \int_0^t \int_{\mathbb{R}} G_\alpha(t-s, x-y) f(s, u_n(s, y)) dy ds \\ &\quad + \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t-s, x-y) g(s, u_n(s-, y), z) \tilde{N}(ds, dy, dz) \end{aligned}$$

It is easy to see that  $u_1 \in \mathcal{E}$ , so by Lemma 3.2, we have for all  $n \geq 1$ ,  $u_n \in \mathcal{E}$ .  
Let

$$F_n(t, x) := E|u_{n+1}(t, x) - u_n(t, x)|^p,$$

and

$$H_n(t) := \sup_{s \leq t} \sup_{x \in \mathbb{R}} F_n(s, x).$$

we have  $\forall t \in [0, T], \forall x \in \mathbb{R}$

$$F_n(t, x) \leq 2^{p-1} [A_n(t, x) + B_n(t, x)] \quad (3.1)$$

with

$$\begin{aligned} A_n(t, x) &:= \\ &E \left( \int_0^t \int_{\mathbb{R}} G_\alpha(t-s, x-y) (f(s, u_n(s, y)) - f(s, u_{n-1}(s, y))) dy ds \right)^p \end{aligned}$$

and

$$\begin{aligned} B_n(t, x) &:= E \left\{ \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t-s, x-y) \right. \\ &\quad \left. (g(s, u_n(s-, y), z) - g(s, u_{n-1}(s-, y), z)) \tilde{N}(ds, dy, dz) \right\}^p. \end{aligned}$$

By condition (H) and Hölder inequality with respect to the measure  $|G_\alpha(t-s, x-y)| ds dy$ , we obtain

$$\begin{aligned} A_n(t, x) &\leq CE \left( \int_0^t \int_{\mathbb{R}} |G_\alpha(t-s, x-y)| |u_n - u_{n-1}|(s, y)^p dy ds \right) \\ &\quad \times \left( \int_0^t \int_{\mathbb{R}} |G_\alpha(t-s, x-y)| dy ds \right)^{p-1}. \\ &\leq C \int_0^t \sup_{y \in \mathbb{R}} E |u_n - u_{n-1}|(s, y)^p \left( \int_{\mathbb{R}} |G_\alpha(t-s, x-y)| dy \right) ds \\ &\leq C \int_0^t H_{n-1}(s) ds. \end{aligned} \quad (3.2)$$

By condition (H), Burkholder inequality and Hölder inequality with respect to the measure  $G_\alpha^2(t-s, x-y)dsdy$  we obtain

$$\begin{aligned}
 B_n(t, x) &\leq CE \left( \int_0^t \int_{\mathbb{R}} G_\alpha^2(t-s, x-y) |u_n - u_{n-1}(s-, y)|^2 dsdy \right)^{\frac{p}{2}} \\
 &\leq C \int_0^t \int_{\mathbb{R}} G_\alpha^2(t-s, x-y) E |u_n - u_{n-1}(s-, y)|^p dsdy \\
 &\quad \times \left( \int_0^t \int_{\mathbb{R}} G_\alpha^2(t-s, x-y) dsdy \right)^{\frac{p}{2}-1} \\
 &\leq C \int_0^t \sup_{y \in \mathbb{R}} |(u_n - u_{n-1})(s, y)|^p \left( \int_{\mathbb{R}} G_\alpha^2(t-s, x-y) dy \right) ds \\
 &\leq C \int_0^t (t-s)^{-\frac{1}{\alpha}} H_{n-1}(s) ds. \tag{3.3}
 \end{aligned}$$

Hence, by (3.1), (3.2), (3.3), we obtain that

$$H_n(t) \leq C \int_0^t (t-s)^{-\frac{1}{\alpha}} H_{n-1}(s) ds$$

for some positive constant  $C$ .

By Lemma 3.3 in [9], the series  $\sum_{n \geq 1} (H_n(t))^{\frac{1}{p}}$  converges uniformly on  $[0, T]$ . Hence the sequence  $(u_n)$  converges in  $L^p(\Omega)$  uniformly on  $[0, T] \times \mathbb{R}$ . Let

$$u(t, x) = L^p(\Omega) - \lim_n u_n(t, x)$$

It is easy to see that  $u \in \mathcal{E}$  and moreover  $u$  satisfies equation (1.2). Thus  $u$  is a solution to equation (1.1) on  $[0, T]$ .

• Proof of the uniqueness: Let  $u$  and  $v$  be two mild solutions of the equation (1.1) on  $[0, T]$  and set

$$H(t) = \sup_{s \leq t} \sup_{x \in \mathbb{R}} E |u(s, x) - v(s, x)|^2.$$

By the same argument as in the proof of existence, we get :

$$H(t) \leq C \int_0^t (t-s)^{-\frac{1}{\alpha}} H(s) ds.$$

Applying Gronwall Lemma, we obtain that  $H(T) = 0$ . This clearly implies that  $u(t, x, \cdot) = v(t, x, \cdot)$  a.s.  $\square$

#### 4. Regularity of the solution

In this section we give the spatial regularity of the solution. This regularity is based on the following lemma proved in [5].

**Lemma 4.1.** *For  $\frac{\alpha+1}{2} < \gamma < \alpha + 1$ , we have*

$$\int_0^{+\infty} \int_{-\infty}^{+\infty} |G_\alpha(v, 1+z) - G_\alpha(v, z)|^\gamma dz dv < \infty.$$

**Theorem 4.2.** *Suppose that  $u_0$  is  $L^p(\Omega)$  bounded for all  $p \geq 2$  and let  $u$  denote the mild solution of the equation (1.1) on  $[0, T]$ , then we have: For  $\alpha < 3$  and for fixed  $t > 0$  the process  $\{u(t, x), x \in \mathbb{R}\}$  has Hölder continuous trajectories with exponent  $\frac{\alpha-1}{2} - \epsilon$ , for any  $\epsilon > 0$ ,  $P$ -a.s.*

*Proof.* The proof is similar to one given by Debbi and Dozzi in [5]. We have

$$\begin{aligned} u(t, x) &= \\ & \int_{\mathbb{R}} G_\alpha(t, x-y) u_0(y) dy + \int_0^t \int_{\mathbb{R}} G_\alpha(t-s, x-y) f(s, u(s, y)) dy ds \\ & \quad + \int_0^{t^+} \int_{\mathbb{R}} \int_{\mathcal{U}} G_\alpha(t-s, x-y) g(s, u(s-, y), z) \tilde{N}(ds, dy, dz) \\ & = u(t, x) + v(t, x) + w(t, x). \end{aligned}$$

It is easy to see that  $u_1$  is smooth function with respect to  $t$  and to  $x$ . By using Lemma 2.2, property (4), we can show that  $x \rightarrow v(t, x)$  is a smooth function, so it is sufficient to estimate the spatial regularity of  $w$ . By Burkholder-Davis-Gundy inequality, condition (H) and the fact the solution is  $L^p(\Omega)$  bounded, we have:

$$\begin{aligned} & E|w(t+h, x) - w(t, x)|^p \\ & \leq C(1 + \sup_{[0, T] \times \mathbb{R}} E|u(t, x)|^p) \times \left\{ \int_0^t \int_{\mathbb{R}} |G_\alpha(v, y+h) - G_\alpha(v, y)|^2 dy dv \right\}^{\frac{p}{2}} \\ & \leq Ch^p \frac{\alpha-1}{2\alpha} \left\{ \int_0^{+\infty} \int_{-\infty}^{+\infty} |G_\alpha(v, 1+z) - G_\alpha(v, z)|^2 dz dv \right\}^{\frac{p}{2}}. \end{aligned}$$

By Lemma 4.1, the last integral converge ( we have taken  $\alpha < 3$  in order that  $\frac{\alpha+1}{2} < 2 < \alpha + 1$  ).

Hence,

$$E|w(t+h, x) - w(t, x)|^p \leq Ch^p \frac{\alpha-1}{2\alpha}$$

we can choose  $p$  as large as we please, so that the result follows from Kolmogorov's Theorem.  $\square$

**Acknowledgment.** I would like to thank the anonymous referee for the careful reading of the first version of the paper and for the valuable suggestions for improving the quality of this article.

## References

- [1] S. Albeverio, J.-L. Wu, and T.-S. Zhang, *Parabolic SPDEs driven by Poisson White Noise*, Stochastic Processes and Their Applications **74** (1998), 21–36.
- [2] E. Saint Loubert Bié, *Etude d'une EDPS conduite par un bruit Poissonien*, Probability Theory and related fields **111** (1998), 287–321.
- [3] R. Dalang and C. Mueller, *Some non-linear s.p.d.e.'s that are second order in time*, Electron. J. Probab. **8** (2003), 1–21. MR MR1961163 (2004a:60118)
- [4] L. Debbi, *On some properties of a High Order fractional differential operator which is not in general selfadjoint*, Applied Mathematical Sciences **1,27** (2007), 1325–1339.
- [5] L. Debbi and M. Dozzi, *On the solutions of nonlinear stochastic fractional partial differential equations in one spatial dimension*, Stoc. Proc. Appl. **115** (2005), 1764–1781.
- [6] N. Fournier, *Malliavin calculus for parabolic SPDEs with jumps*, Stochastic Processes and Their Applications **87** (2000), 115–147.
- [7] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, North-Holland Publishing Company. Mathematical Library 24., Holland, 1989.
- [8] I. Podlubny, *Fractional Differential equations: an Introduction to Fractional Derivatives, Fractional Differential equations, to Methods of Their Solution and Some of their Applications*, Academic Press, San Diego, CA., 1999.
- [9] J.B. Walsh., *An Introduction to stochastic partial differential equations*, Lecture Notes in Mathematics 1180, Springer Berlin / Heidelberg, 1986, pp. 266–437.

FRACTIONAL SPDES DRIVEN BY POISSON WHITE NOISE

- [10] J. Zabczyk., *Symmetric solutions of semilinear stochastic equations*, Lecture Notes in Mathematics 1390, Springer Berlin / Heidelberg, 1988, pp. 237–256.

SALAH HAJJI  
Department of Mathematics  
Faculty of Sciences Semlalia  
Cadi Ayyad University  
BP. 2390  
Marrakesh, MOROCCO.  
s.hajji@ucam.ac.ma