

BERTIN DIARRA

***p*-adic Clifford algebras**

Annales mathématiques Blaise Pascal, tome 1, n° 1 (1994), p. 85-103

http://www.numdam.org/item?id=AMBP_1994__1_1_85_0

© Annales mathématiques Blaise Pascal, 1994, tous droits réservés.

L'accès aux archives de la revue « Annales mathématiques Blaise Pascal » (<http://math.univ-bpclermont.fr/ambp/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

P-ADIC CLIFFORD ALGEBRAS

Bertin DIARRA

In a previous paper [2], we gave the index of the standard quadratic form of rank n over the field of p -adic numbers. Here, we recover, as a consequence, the structure of the associated Clifford algebra.

The classification of all (equivalence classes of) quadratic forms over a p -adic field is well known (cf.[5]), with this classification, one is able to classify all p -adic Clifford algebras.

I - INTRODUCTION

Let K be a field of characteristic $\neq 2$ and E a vector space over K of finite dimension n . A mapping $q : E \rightarrow K$ is a *quadratic form* over E if there exists a bilinear symmetric form $f : E \times E \rightarrow K$ such that

$$q(x) = f(x, x) \quad \text{and} \quad f(x, y) = \frac{1}{2}[q(x+y) - q(x) - q(y)]$$

We assume that q is *regular*, that is f is non-degenerated.

An element $x \in E$ is *isotropic* if $q(x) = 0$. Let V be a subspace of E ; the orthogonal subspace of V is the set $V^\perp = \{y \in E / f(x, y) = 0 \text{ for all } x \in V\}$. The subspace V is called *totally isotropic* if $V \subset V^\perp$. It is well known (cf. for example [1]) that any totally isotropic subspace is contained in a maximal totally isotropic subspace. The maximal totally isotropic subspaces have the same dimension ν , called the *index* of q and $2\nu \leq n$. If $2\nu = n$, then (E, q) is called a *hyperbolic* space and for the case $n = 2$, one says hyperbolic plane. The index $\nu = 0$ iff $q(x) \neq 0$ for $x \neq 0$ i.e. (E, q) is *anisotropic*.

Let $E = K^n$ and $B = (e_1, \dots, e_n)$ be the canonical basis of E ; the standard quadratic form q_0 is the quadratic form associated to the bilinear form

$$\langle x, y \rangle = \sum_{j=1}^n x_j y_j \quad ; \quad \text{where } x = \sum_{j=1}^n x_j e_j \quad \text{and} \quad y = \sum_{j=1}^n y_j e_j \quad ;$$

hence $q_0(x) = \langle x, x \rangle = \sum_{j=1}^n x_j^2$.

Let (E, q) be a quadratic space, possibly non regular ; an algebra $C = C(E, q)$ over K , with unit 1, is said to be a *Clifford algebra* for (E, q) if

(i) There exists a one-to-one linear mapping $\rho : E \rightarrow C$ such that $\rho(x)^2 = q(x) \cdot 1$.

(ii) For every algebra A with unit 1 and linear mapping $\phi : E \rightarrow A$ satisfying $\phi(x)^2 = q(x) \cdot 1$, there exists an algebra homomorphism $\tilde{\phi} : C \rightarrow A$ such that $\tilde{\phi} \circ \rho = \phi$.

Clifford algebra exists and is unique up algebra isomorphism (cf. for instance [1] or [3]). For example, let $K \langle X_1, \dots, X_n \rangle$ be the free algebra with free system of generators X_1, \dots, X_n and I be the two-sided ideal of $K \langle X_1, \dots, X_n \rangle$ generated by $X_i X_j + X_j X_i - 2f(e_i, e_j) \cdot 1, 1 \leq i, j \leq n$, where (e_1, \dots, e_n) is an orthogonal basis of (E, q) ; then $C(E, q) = K \langle X_1, \dots, X_n \rangle / I$.

II - THE P-ADIC STANDARD QUADRATIC FORM q_0

II - 1 . The index of q_0

Let p be a prime number and \mathbf{Q}_p be the p -adic field i.e. the completion of the field of rational numbers \mathbf{Q} for the p -adic absolute value.

We denote by $[\alpha]$ the integral part of the real number α .

Proposition 1 [2]

The standard quadratic form $q_0(x) = \sum_{j=1}^n x_j^2$ over $E = \mathbf{Q}_p^n$ has index

(i) $\nu = \left[\frac{n}{2} \right]$ if $p \equiv 1 \pmod{4}$

(ii) $\nu = \left[\frac{n}{2} \right]$ if $p \equiv 3 \pmod{4}$ and $n \not\equiv 2 \pmod{4}$

(iii) $\nu = \left[\frac{n}{2} \right] - 1$ if $p \equiv 3 \pmod{4}$ and $n \equiv 2 \pmod{4}$

Proof :

1°) If $p \equiv 1 \pmod{4}$, it is well known that $i = \sqrt{-1} \in \mathbf{Q}_p$. Let $\nu = \left[\frac{n}{2} \right]$ and $\epsilon_j = i e_{2j-1} + e_{2j}$, $1 \leq j \leq \nu$, then $V = \bigoplus_{j=1}^{\nu} \mathbf{Q}_p \epsilon_j$ is a maximal totally isotropic subspace of $E = \mathbf{Q}_p^n$.

2°) $p \equiv 3 \pmod{4}$

Therefore $i \notin \mathbf{Q}_p$ and if $n = 2$ the index of q_0 is 0.

If $n = 3$, applying Chevalley's theorem and Newton's method to $q_0(x) = x_1^2 + x_2^2 + x_3^2$ we find $a, b \in \mathbf{Q}_p$, $a \neq 0, b \neq 0$, such that $a^2 + b^2 + 1 = 0$. Therefore $\epsilon_1 = a e_1 + b e_2 + e_3$ is isotropic in \mathbf{Q}_p^3 and $\nu = \left[\frac{3}{2} \right] = 1$.

(a) For $n = 4m$, put $\epsilon_{2j-1} = a e_{4j-3} + b e_{4j-2} + e_{4j-1}$ and $\epsilon_{2j} = -b e_{4j-3} + a e_{4j-2} + e_{4j}$, $1 \leq j \leq m$. It is clear that $q_0(\epsilon_{2j-1}) = q_0(\epsilon_{2j}) = a^2 + b^2 + 1 = 0$ and $\langle \epsilon_{2j-1}, \epsilon_{2j} \rangle = -ab + ab = 0$. Therefore $V = \bigoplus_{j=1}^m (\mathbf{Q}_p \epsilon_{2j-1} \oplus \mathbf{Q}_p \epsilon_{2j})$ is a totally isotropic subspace of \mathbf{Q}_p^n and $\nu = 2m = \left[\frac{n}{2} \right]$.

If $n = 4m + 1$, with the same notations as above the subspace V is totally isotropic in \mathbf{Q}_p^n and $\nu = 2m = \left[\frac{n}{2} \right]$.

On the other hand if $n = 4m + 3$ the subspaces $V = \bigoplus_{j=1}^m (\mathbf{Q}_p \epsilon_{2j-1} \oplus \mathbf{Q}_p \epsilon_{2j})$ and $\mathbf{Q}_p \epsilon_{2m+1}$ where $\epsilon_{2m+1} = a e_{4m+1} + b e_{4m+2} + e_{4m+3}$, are totally isotropic and orthogonal. Therefore $V_0 = V \oplus \mathbf{Q}_p \epsilon_{2m+1}$ is totally isotropic and $\nu = 2m + 1 = \left[\frac{n}{2} \right]$.

(b) If $n = 4m + 2$, let $V = \bigoplus_{j=1}^m (\mathbf{Q}_p \epsilon_{2j-1} \oplus \mathbf{Q}_p \epsilon_{2j})$ be as above. It is easy to verify that if $x \in \mathbf{Q}_p^n$ is isotropic and x is orthogonal to V then $x \in V$. Therefore V is a maximal totally isotropic subspace of \mathbf{Q}_p^n and $\nu = 2m = \left[\frac{n}{2} \right] - 1$.

Proposition 2 : Let $p = 2$.

Let $n = 8m + s$, $0 \leq s \leq 7$.

The standard quadratic form $q_0(x) = \sum_{j=1}^n x_j^2$ over $E = \mathbf{Q}_2^n$ has index

$$(i) \quad \nu = 4m \quad \text{if} \quad 0 \leq s \leq 4$$

$$(ii) \quad \nu = 4m + t \quad \text{if} \quad s = 4 + t, \quad 1 \leq t \leq 3$$

Proof :

1°) If $1 \leq n \leq 4$, then the index of q_0 is 0.

Indeed, this is clear when $n = 1$.

If $n = 2$, let $x = x_1 e_1 + x_2 e_2 \in \mathbf{Q}_2^2$ be isotropic and different from 0 i.e. $q_0(x) = x_1^2 + x_2^2 = 0$ and say $x_2 \neq 0$. Therefore $1 + a^2 = 0$ with $a = x_1 x_2^{-1}$ and $v_2(a) = 0$ i.e. $a = 1 + 2^\mu a_0$, $\mu \geq 1$, $v_2(a_0) = 0$.

Then $1 + a^2 = 2 + 2^{\mu+1} a_0 + 2^{2\mu} a_0^2 = 0$ or $1 + 2^\mu a_0 + 2^{2\mu-1} a_0 = 0$; in other words $1 \equiv 0 \pmod{2}$; a contradiction.

In the same way, one shows that if $n = 3$ or 4 , the index of q_0 is 0.

2°) $n = 5$

Let $x_0 = 2e_1 + e_2 + e_3 + e_4 + e_5 \in \mathbf{Q}_2^5$, then $q_0(x_0) = 8$ and $\frac{\partial q_0}{\partial x_j}(x_0) = 2 \not\equiv 0 \pmod{4}$,

$2 \leq j \leq 5$

By Newton's method there exists

$x = \sum_{j=1}^5 x_j e_j \in \mathbf{Q}_2^5$ such that $q_0(x) = 0$ with $x_1 \equiv 2 \pmod{8}$, $x_j \equiv 1 \pmod{8}$, $2 \leq j \leq 5$.

Put $a = x_1 x_5^{-1}$, $b = x_2 x_5^{-1}$, $c = x_3 x_5^{-1}$, $d = x_4 x_5^{-1}$, then $a^2 + b^2 + c^2 + d^2 + 1 = 0$.

The two following elements of \mathbf{Q}_2^5

$$e_1 = a e_1 + b e_2 + c e_3 + d e_4 + e_5$$

$$e'_1 = -a e_1 - b e_2 - c e_3 - d e_4 + e_5$$

are isotropic with $\langle e_1, e'_1 \rangle = 2$. Hence $H = \mathbf{Q}_2 e_1 \oplus \mathbf{Q}_2 e'_1$ is a hyperbolic plane in \mathbf{Q}_2^5 . Let

$U = H^\perp$ be the orthogonal subspace of H in \mathbf{Q}_2^5 . The following three elements of \mathbf{Q}_2^5 :

$$u_1 = b e_1 - a e_2 + d e_3 - c e_4$$

$$u_2 = e_1 - \frac{ac + bd}{c^2 + d^2} e_3 + \frac{bc - ad}{c^2 + d^2} e_4$$

$$u_3 = e_2 + \frac{ad - bc}{c^2 + d^2} e_3 - \frac{ac + bd}{c^2 + d^2} e_4$$

are elements of U , with

$$q_0(u_1) = -1, \quad q_0(u_2) = -\frac{1}{c^2 + d^2} = q_0(u_3)$$

Furthermore $\langle u_i, u_j \rangle = 0$ if $1 \leq i \neq j \leq 3$, and (u_1, u_2, u_3) is a basis of U .

For every $u = y_1 u_1 + y_2 u_2 + y_3 u_3 \in U$ we have $q_0(u) = y_1^2 q_0(u_1) + y_2^2 q_0(u_2) + y_3^2 q_0(u_3) = -\frac{c^2 y_1^2 + d^2 y_2^2 + y_3^2}{c^2 + d^2}$ and $q_0(u) = 0$ iff $u = 0$ because the standard quadratic form of rank 4 is anisotropic. In other words (U, q_0) is anisotropic and $\mathbf{Q}_2^5 = H \perp U$ is a Witt decomposition of (\mathbf{Q}_2^5, q_0) . Hence the index of q_0 is 1.

3°) $\underline{n} = 8m + s, \quad 0 \leq s \leq 4.$

Put, for $0 \leq j \leq m - 1$

$$(1) \quad \begin{cases} \epsilon_{j,1} = a e_{8j+1} + b e_{8j+2} + c e_{8j+3} + d e_{8j+4} + e_{8j+5} \\ \epsilon_{j,2} = -b e_{8j+1} + a e_{8j+2} + d e_{8j+3} - c e_{8j+4} + e_{8j+6} \\ \epsilon_{j,3} = -d e_{8j+1} + c e_{8j+2} - b e_{8j+3} + a e_{8j+4} + e_{8j+7} \\ \epsilon_{j,4} = c e_{8j+1} + d e_{8j+2} - a e_{8j+3} - b e_{8j+4} + e_{8j+8} \end{cases}$$

and

$$(2) \quad \begin{cases} \epsilon'_{j,1} = -a e_{8j+1} - b e_{8j+2} - c e_{8j+3} - d e_{8j+4} + e_{8j+5} \\ \epsilon'_{j,2} = b e_{8j+1} - a e_{8j+2} - d e_{8j+3} + c e_{8j+4} + e_{8j+6} \\ \epsilon'_{j,3} = d e_{8j+1} - c e_{8j+2} + b e_{8j+3} - a e_{8j+4} + e_{8j+7} \\ \epsilon'_{j,4} = -c e_{8j+1} - d e_{8j+2} + a e_{8j+3} + b e_{8j+4} + e_{8j+8} \end{cases}$$

A straightforward computation shows that $\langle \epsilon_{i,k}, \epsilon_{j,l} \rangle = 0 = \langle \epsilon'_{i,k}, \epsilon'_{j,l} \rangle, \quad 0 \leq i, j \leq m - 1; \quad 1 \leq k, l \leq 4$ and $\langle \epsilon_{j,l}, \epsilon'_{j,l} \rangle = 2; \quad 0 \leq j \leq m - 1; \quad 1 \leq l \leq 4.$ Furthermore $\langle \epsilon_{i,k}, \epsilon'_{j,l} \rangle = 0$ if $(i, k) \neq (j, l).$

Hence the subspaces $V = \bigoplus_{\substack{j=0 \\ 1 \leq l \leq 4}}^{m-1} \mathbf{Q}_2 \epsilon_{j,l}$ and $W = \bigoplus_{\substack{j=0 \\ 1 \leq l \leq 4}}^{m-1} \mathbf{Q}_2 \epsilon'_{j,l}$ are isotropic with

$$V \cap W = (0)$$

Therefore $H = V \oplus W$ is a hyperbolic subspace of $E = \mathbf{Q}_2^{8m+s}$, with $\dim V = \dim W = 4m.$

But $E = E_m \perp E_s$ (orthogonal sum) where $E_m = \bigoplus_{j=1}^{8m} \mathbf{Q}_2 e_j$ and $E_s = \bigoplus_{k=1}^s \mathbf{Q}_2 e_{8m+k} \simeq$

\mathbf{Q}_2^s .

If $s = 0$, we have $E = E_m = V \oplus W = H$ and (E, q_0) is a hyperbolic space with index $4m$.

If $1 \leq s \leq 4$; $E = E_m \perp E_s$ with $E_m = V \oplus W = H$. Since $1 \leq \dim E_s = s \leq 4$, the standard quadratic space (E_s, q_0) is anisotropic. Consequently $E = (V \oplus W) \perp E_s$ is a Witt decomposition of E and the index of q_0 is $4m$.

4°) $\underline{n = 8m + 4 + t}$, $1 \leq t \leq 3$.

a) $\underline{n = 8m + 5}$

With the same notations as above, we have $E = E_m \perp E_5$ where $E_5 = \bigoplus_{k=1}^5 \mathbf{Q}_2 e_{8m+k} \simeq$

\mathbf{Q}_2^5 .

Let us write, as for $n = 5$,

$$(3) \quad \begin{cases} \epsilon_{4m+1} &= a e_{8m+1} + b e_{8m+2} + c e_{8m+3} + d e_{8m+4} + e_{8m+5} \\ \epsilon'_{4m+1} &= -a e_{8m+1} - b e_{8m+2} - c e_{8m+3} - d e_{8m+4} + e_{8m+5} \end{cases}$$

and

$$(4) \quad \begin{cases} u_{m+1} &= b e_{8m+1} - a e_{8m+2} + d e_{8m+3} - c e_{8m+4} \\ u_{m+2} &= e_{8m+1} - \frac{ac+bd}{c^2+d^2} e_{8m+3} + \frac{bc-ad}{c^2+d^2} e_{8m+4} \\ u_{m+3} &= e_{8m+2} + \frac{ad-bc}{c^2+d^2} e_{8m+3} - \frac{ac+bd}{c^2+d^2} e_{8m+4} \end{cases}$$

The subspace $U_5 = \bigoplus_{h=1}^3 \mathbf{Q}_2 u_{m+h}$ of E_5 is anisotropic. On the other hand, $q_0(\epsilon_{4m+1}) =$

$0 = q_0(\epsilon'_{4m+1})$; $\langle \epsilon_{4m+1}, \epsilon'_{4m+1} \rangle = 2$ and $\epsilon_{4m+1}, \epsilon'_{4m+1}$ are orthogonal to U_5 . Therefore $V_0 = V \oplus \mathbf{Q}_2 \epsilon_{4m+1}$ and $W_0 = W \oplus \mathbf{Q}_2 \epsilon'_{4m+1}$ are isotropic subspaces of E and $E = (V_0 \oplus W_0) \perp U_5$ is a Witt decomposition of E . Hence the index of q_0 is $\dim V_0 = \dim W_0 = 4m+1$.

(b) $\underline{n = 8m+6}$.

As before, we have $E = E_m \perp E_6$ where $E_6 = \bigoplus_{k=1}^6 \mathbf{Q}_2 e_{8m+k} \supset E_5$; hence ϵ_{4m+1} and

$\epsilon'_{4m+1} \in E_6$.

Let us put

$$(5) \quad \begin{cases} \epsilon_{4m+2} &= -b e_{8m+1} + a e_{8m+2} + d e_{8m+3} - c e_{8m+4} + e_{8m+6} \\ \epsilon'_{4m+2} &= b e_{8m+1} - a e_{8m+2} - d e_{8m+3} + c e_{8m+4} + e_{8m+6} \end{cases}$$

and

$$(6) \quad \begin{cases} \omega_{m+1} &= e_{8m+1} + \frac{bd-ac}{c^2+d^2} e_{8m+3} - \frac{ad+bc}{c^2+d^2} e_{8m+4} \\ \omega_{m+2} &= e_{8m+2} - \frac{bc+ad}{c^2+d^2} e_{8m+3} + \frac{ac-bd}{c^2+d^2} e_{8m+4} \end{cases}$$

The subspace $U_6 = \mathbb{Q}_2 \omega_{m+1} \oplus \mathbb{Q}_2 \omega_{m+2}$ of E_6 is anisotropic. Moreover, $q_0(\epsilon_{4m+2}) = 0 = q_0(\epsilon'_{4m+2})$; $\langle \epsilon_{4m+2}, \epsilon'_{4m+2} \rangle = 2$ and $\epsilon_{4m+2}, \epsilon'_{4m+2}$ are orthogonal to U_6 . Therefore $V_1 = V_0 \oplus \mathbb{Q}_2 \epsilon_{4m+2}$ and $W_1 = W_0 \oplus \mathbb{Q}_2 \epsilon'_{4m+2}$ are isotropic subspaces of E and $E = (V_1 \oplus W_1) \perp U_6$ is a Witt decomposition of E . Hence the index of q_0 is $\dim V_1 = \dim W_1 = 4m + 2$.

(c) $\quad \underline{n = 8m+7.}$

We have $E = E_m \perp E_7$, where $E_7 = \bigoplus_{k=1}^7 \mathbb{Q}_2 e_{8m+k} \supset E_6$.

Let us write

$$(7) \quad \begin{cases} \epsilon_{4m+3} &= -d e_{8m+1} + c e_{8m+2} - b e_{8m+3} + a e_{8m+4} + e_{8m+7} \\ \epsilon'_{4m+3} &= d e_{8m+1} - c e_{8m+2} + b e_{8m+3} - a e_{8m+4} + e_{8m+7} \end{cases}$$

and

$$(8) \quad u_m = c e_{8m+1} + d e_{8m+2} - a e_{8m+3} - b e_{8m+4}$$

The subspace $U_7 = \mathbb{Q}_2 u_m$ of E_7 is anisotropic. Furthermore $q_0(\epsilon_{4m+3}) = 0 = q_0(\epsilon'_{4m+3})$; $\langle \epsilon_{4m+3}, \epsilon'_{4m+3} \rangle = 2$ and $\epsilon_{4m+3}, \epsilon'_{4m+3}$ are orthogonal to U_7 . Therefore $V_2 = V_1 \oplus \mathbb{Q}_2 \epsilon_{4m+3}$ and $W_2 = W_1 \oplus \mathbb{Q}_2 \epsilon'_{4m+3}$ are isotropic subspaces of E and $E = (V_2 \oplus W_2) \perp U_7$ is a Witt decomposition of E . Hence the index of q_0 is $\dim V_2 = \dim W_2 = 4m + 3$.

Remark

Let K be a non formally real field. The level of K is the least integer s such that

$$-1 = \sum_{j=1}^s a_j^2 \text{ where } a_j \in K, a_j \neq 0. \text{ It is well known that } s = 2^r, r \geq 0 \text{ (c f. [3] or [4]).}$$

The level of a p -adic field is 1 if $p \equiv 1 \pmod{4}$; 2 if $p \equiv 3 \pmod{4}$ and 4 if $p = 2$.

If the level of a field K is 1 (resp. 2, resp.4) then the index of the standard quadratic form over K^n is given by Proposition 1 - (i) [resp. Prop.1 - (ii) - (iii) , resp. Prop.2].

More generally let K be a field of level $s = 2^r, r \geq 0$. If we write for any integer $n, n = m2^{r+1} + a$ where $0 \leq a \leq 2^{r+1} - 1$; then the index of the standard quadratic form over K^n is

$$(i) \quad \nu = m2^r \quad \text{if } 0 \leq a \leq 2^r$$

$$(ii) \quad \nu = m2^r + t \quad \text{if } a = 2^r + t, \quad 1 \leq t \leq 2^r - 1.$$

II - 2 The Clifford algebra $C(\mathbb{Q}_p^n, q_0)$

The following results can be deduced from a general setting (cf. [3] p. 128-129). Here we establish them by using the computation of the index of q_0 made in II-1.

Let us recall that if E is a vector space over a field K then the exterior algebra $\wedge(E)$ is the Clifford algebra associated to the null quadratic form over E .

On the other hand , let (E, q) be a regular quadratic space over K . If $E = V \oplus W$ is a hyperbolic space (V and W being maximal totally isotropic subspaces) , it is well known that the Clifford algebra $C(E, q)$ is isomorphic to $End(\wedge(V))$, the space of linear endomorphisms of the vector space $\wedge(V)$. Furthermore the subalgebra of the even elements of $C(E, q)$, say $C_+(E, q)$ is isomorphic to $End(\wedge_+(V)) \times End(\wedge_-(V))$ where $\wedge_+(V)$ (resp. $\wedge_-(V)$) is the subspace of the even (resp. odd) elements of $\wedge(V)$.

Generally, if $E = (V \oplus W) \perp U$ is a Witt decomposition of E , then $C(E, q) \simeq End(\wedge(V)) \otimes_2 C(U, q)$, the tensor product of $\mathbb{Z}/2\mathbb{Z}$ - graded algebras (cf. for example [1]).

If $dim E = n$, then $dim C(E, q) = 2^n = dim \wedge(E)$.

If $a, b \in K^*$, we denote by $\left(\frac{a, b}{K}\right)$ the associated quaternion algebra : i.e. the algebra generated by i, j with $i^2 = a$; $j^2 = b$; $ij = -ji$. Also $\left(\frac{a, b}{K}\right)$ is the Clifford algebra of the rank 2 quadratic form $q(x) = ax_1^2 + bx_2^2$.

Let us write $M(n, K)$ the algebra of the $n \times n$ matrices with coefficients in K .

Theorem 1 : $p \equiv 1 \pmod{4}$

- (i) If $n = 2m$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^m, \mathbf{Q}_p)$
- (ii) If $n = 2m + 1$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^m, \mathbf{Q}_p) \oplus M(2^m, \mathbf{Q}_p)$

Proof

Indeed, if $n = 2m$, then (\mathbf{Q}_p^n, q_0) is a hyperbolic space.

It follows that $C(\mathbf{Q}_p^n, q_0) \simeq \text{End}(\wedge(\mathbf{Q}_p^m))$.

And, if $n = 2m + 1$, we have a Witt decomposition $\mathbf{Q}_p^n = (V \oplus W) \perp U$ where $U = \mathbf{Q}_p e_n$.

It follows that $C(U, q_0) \simeq \mathbf{Q}_p \oplus \mathbf{Q}_p$ which gives (ii)

Theorem 2 : $p \equiv 3 \pmod{4}$

- (i) If $n = 4m$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m}, \mathbf{Q}_p)$
- (ii) If $n = 4m + 1$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m}, \mathbf{Q}_p) \oplus M(2^{2m}, \mathbf{Q}_p)$
- (iii) If $n = 4m + 2$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m+1}, \mathbf{Q}_p)$
- (iv) If $n = 4m + 3$, then $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m+1}, \mathbf{Q}_p[i])$

with $i = \sqrt{-1}$.

Proof :

The case (i) is evident, since \mathbf{Q}_p^{2m} is a hyperbolic space.

If $n = 4m + 1$, we have a Witt decomposition $\mathbf{Q}_p^n = (V \oplus W) \perp U$ where $U = \mathbf{Q}_p u$ with $u = a e_{4m-3} + b e_{4m-2} + e_{4m-1} - e_{4m+1}$ and $q_0(u) = a^2 + b^2 + 1 + 1 = 1$. It follows that $C(U, q_0) \simeq \mathbf{Q}_p \oplus \mathbf{Q}_p$, which gives (ii).

If $n = 4m + 2$, we have a Witt decomposition $\mathbf{Q}_p^n = (V \oplus W) \perp U$ where $U = \mathbf{Q}_p u_1 \oplus \mathbf{Q}_p u_2$ and $u_1 = a e_{4m-3} + b e_{4m-2} + e_{4m-1} + a e_{4m+1} + b e_{4m+2}$
 $u_2 = -b e_{4m-3} + a e_{4m-2} + e_{4m} - b e_{4m+1} + a e_{4m+2}$

Furthermore $\langle u_1, u_2 \rangle = 0$, $q_0(u_1) = -1 = q_0(u_2)$ and $C(U, q_0) \simeq \left(\frac{-1, -1}{\mathbf{Q}_p} \right)$. This quaternion algebra contains an element z with $N(z) = a^2 + b^2 + 1 = 0$. Hence $\left(\frac{-1, -1}{\mathbf{Q}_p} \right) \simeq M(2, \mathbf{Q}_p)$ and finally we have $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m}, \mathbf{Q}_p) \otimes_2 M(2, \mathbf{Q}_p) \simeq M(2^{2m+1}, \mathbf{Q}_p)$.

If $n = 4m + 3$, we have a Witt decomposition $\mathbf{Q}_p^n = (V \oplus W) \perp U$ where $U = \mathbf{Q}_p u$, with $u = -b e_{4m+1} + a e_{4m+2}$ and $q_0(u) = b^2 + a^2 = -1$. Hence $C(U, q_0) \simeq \mathbf{Q}_p[i]$, because $u^2 = q_0(u) = -1$.

We conclude that $C(\mathbf{Q}_p^n, q_0) \simeq M(2^{2m+1}, \mathbf{Q}_p[i])$.

In the proof of the foregoing theorem, one needs the following lemma

Lemma :

Let K be a field (char. $\neq 2$), $c, d \in K^*$ such that $c^2 + d^2 \neq 0$.

If $\sigma = \frac{1}{c^2 + d^2}$, then $\left(\frac{-\sigma, -\sigma}{K}\right) \simeq \left(\frac{-1, -1}{K}\right)$.

If the two-rank quadratic forms $q_1(x) = -\sigma x_1^2 - \sigma x_2^2$ and $q_2(x) = -x_1^2 - x_2^2$ are equivalent, then their Clifford algebras are isomorphic. But, putting $x_1 = cx'_1 + dx'_2$ and $x_2 = dx'_1 - cx'_2$, we have $q_1(u(x')) = -\sigma(cx'_1 + dx'_2)^2 - \sigma(dx'_1 - cx'_2)^2 = -\sigma(c^2 + d^2)(x_1'^2 + x_2'^2) = q_2(x')$. Hence q_1 and q_2 are equivalent and the lemma is proved.

Remark

The quaternion algebra $\left(\frac{-1, -1}{\mathbf{Q}_2}\right) = \mathbf{H}_2$ is a skew field.

Indeed, for any $z \in \mathbf{H}_2 = \left(\frac{-1, -1}{\mathbf{Q}_2}\right)$, $z \neq 0$, the norm of z is $N(z) = x_0^2 + x_1^2 + x_2^2 + x_3^2 \neq 0$ (the standard quadratic form of rank 4 over \mathbf{Q}_2 is anisotropic).

Theorem 3 : $p = 2$

The Clifford algebra $C(\mathbf{Q}_2^n, q_0)$ is isomorphic to :

- (0) $End(\wedge(\mathbf{Q}_2^{4m})) \simeq M(2^{4m}, \mathbf{Q}_2)$, if $n = 8m$
- (1) $M(2^{4m}, \mathbf{Q}_2) \oplus M(2^{4m}, \mathbf{Q}_2)$, if $n = 8m + 1$
- (2) $M(2^{4m+1}, \mathbf{Q}_2)$, if $n = 8m + 2$
- (3) $M(2^{4m+1}, \mathbf{Q}_2[i])$, with $i = \sqrt{-1}$, if $n = 8m + 3$
- (4) $M(2^{4m+1}, \mathbf{H}_2)$, if $n = 8m + 4$
- (5) $M(2^{4m+1}, \mathbf{H}_2) \oplus M(2^{4m+1}, \mathbf{H}_2)$, if $n = 8m + 5$
- (6) $M(2^{4m+2}, \mathbf{H}_2)$, if $n = 8m + 6$
- (7) $M(2^{4m+3}, \mathbf{Q}_2[i])$, if $n = 8m + 7$

Proof

According to the proof of Proposition 2, if $n = 8m + s, 0 \leq s \leq 7$, then $\mathbf{Q}_2^n = (V \oplus W) \perp E_s$, where V and W are totally isotropic subspaces of dimension $4m$, and $(E_s, q_0) \simeq$

(\mathbf{Q}_2^s, q_0) . It follows that $C(\mathbf{Q}_2^n, q_0) \simeq \text{End}(\wedge(\mathbf{Q}_p^{4m})) \otimes_2 C(\mathbf{Q}_2^s, q_0)$. It is easy to see that $C(\mathbf{Q}_2, q_0) \simeq \mathbf{Q}_2 \oplus \mathbf{Q}_2$; $C(\mathbf{Q}_2^2, q_0) \simeq \begin{pmatrix} 1, 1 \\ \mathbf{Q}_2 \end{pmatrix} \simeq M(2, \mathbf{Q}_2)$ and $C(\mathbf{Q}_2^3, q_0) \simeq M(2, \mathbf{Q}_2[i])$.

If $s = 4$, the subalgebra, generated by e_1e_2, e_2e_4 and e_1e_4 , is isomorphic to $\begin{pmatrix} -1, -1 \\ \mathbf{Q}_2 \end{pmatrix} = \mathbf{H}_2$. Hence $C(\mathbf{Q}_2^4, q_0) \simeq M(2, \mathbf{H}_2)$.

If $s = 5$, then $\mathbf{Q}_2^5 = F \perp U$, where F is a hyperbolic plane and U a three-dimensional anisotropic subspace, with orthogonal basis (u_1, u_2, u_3) satisfying $q_0(u_1) = -1, q_0(u_2) = -\sigma = q_0(u_3)$. ($\sigma = \frac{1}{c^2+d^2}$ and $a, b, c, d \in \mathbf{Q}_2$ such that $a^2 + b^2 + c^2 + d^2 + 1 = 0$).

Therefore $C_+(U, q_0) \simeq \begin{pmatrix} -\sigma, -\sigma \\ \mathbf{Q}_2 \end{pmatrix} \simeq \mathbf{H}_2$; C_+ stands for the even subalgebra. But in $C(U, q_0), (u_1u_2u_3)^2 = \sigma^2$ is a square in \mathbf{Q}_2 ; therefore $C(U, q_0) \simeq \mathbf{H}_2 \oplus \mathbf{H}_2$. Furthermore $C(\mathbf{Q}_2^5, q_0) \simeq C(F, q_0) \otimes_2 C(U, q_0) \simeq M(2, \mathbf{H}_2) \oplus M(2, \mathbf{H}_2)$, because $C(F, q_0) \simeq M(2, \mathbf{Q}_2)$.

If $s = 6$, then $\mathbf{Q}_2^6 = F \perp U$, where F is a hyperbolic space of dimension 4 and U a two-dimensional anisotropic subspace with an orthogonal basis (u_1, u_2) satisfying $q(u_1) = -\sigma = q(u_2)$. Therefore $C(U, q_0) \simeq \begin{pmatrix} -\sigma, -\sigma \\ \mathbf{Q}_2 \end{pmatrix} \simeq \mathbf{H}_2$. And consequently $C(\mathbf{Q}_2^6, q_0) \simeq C(F, q_0) \otimes_2 C(U, q_0) \simeq M(2^2, \mathbf{H}_2)$.

If $s = 7$, then $\mathbf{Q}_2^7 = F \perp U$, where F is a hyperbolic space of dimension 6 and $U = \mathbf{Q}_2u$, with $q_0(u) = -1$. Hence $C(U, q_0) \simeq \mathbf{Q}_2[i]$ and $C(\mathbf{Q}_2^7, q_0) \simeq M(2^3, \mathbf{Q}_2[i])$.

One deduces the isomorphisms of the theorem from $C(\mathbf{Q}_2^n, q_0) \simeq M(2^{4m}, \mathbf{Q}_2) \otimes_2 C(\mathbf{Q}_2^s, q_0)$.

N.B : A classical way to prove the above theorems is based on the isomorphisms

$$C(K^{n+2}, q_0) \simeq C(K^n, -q_0) \otimes C(K^2, q_0)$$

and $C(K^{n+2}, -q_0) \simeq C(K^n, q_0) \otimes C(K^2, -q_0)$

which give first 8-periodicity, etc ...

($-q_0$ is the opposite of the standard quadratic form q_0)

III - THE FAMILIES OF P-ADIC CLIFFORD ALGEBRAS

III-1. Equivalent classes of the p-adic quadratic forms

Let $a, b \in \mathbf{Q}_p^* = \mathbf{Q}_p \setminus \{0\}$. The Hilbert symbol (a, b) is defined by $(a, b) = 1$ if the quadratic form of rank 3, $q'(x) = x_0^2 - ax_1^2 - bx_2^2$ is isotropic $(a, b) = -1$ otherwise.

N.B. $(a, b) = 1$ iff $\left(\frac{a, b}{\mathbf{Q}_p}\right) \simeq M(2, \mathbf{Q}_p)$.

Let E be a vector space over \mathbf{Q}_p of dimension n . Let us consider a regular quadratic form q over E . If $(e_j)_{1 \leq j \leq n}$ is an orthogonal basis of E and $a_j = q(e_j)$; then the discriminant $d(q)$ of q is equal to $a_1 \dots a_n$ in the group $M_p = \mathbf{Q}_p^*/\mathbf{Q}_p^2$. Let $\epsilon(q) = \prod_{1 \leq i < j \leq n} (a_i, a_j)$.

Theorem A

(i) *The p-adic regular quadratic forms q and q' of rank n are equivalent iff $d(q) = d(q')$ and $\epsilon(q) = \epsilon(q')$.*

(ii) *Let $d \in M_p$ and $\epsilon = \pm 1$. There exists a p-adic regular quadratic form q such that $d(q) = d$ and $\epsilon(q) = \epsilon$ iff*

- (a) $n = 1$ and $\epsilon = 1$
- (b) $n = 2$ and $(d, \epsilon) \neq (-1, -1)$
- (c) $n \geq 3$

Proof : cf. [5]

According to that proof of Theorem A, one can give, explicitly, representatives of the equivalence classes of p-adic regular quadratic forms.

Let us recall that $M_2 = \{\pm 1, \pm 2, \pm 5, \pm 10\}$ and $M_p = \{1, p, \omega, \omega p\}$ if $p \neq 2$, where ω is a unit such that $\left(\frac{\omega}{p}\right) = -1$; $\left(\frac{-}{p}\right)$ = the Legendre symbol. Furthermore $-1 = 1$ in M_p if $p \equiv 1 \pmod{4}$ and $M_p = \{1, p, -1, -p\}$ if $p \equiv 3 \pmod{4}$.

We are content ourselves here, with the primes p different from 2. Then a complete set of representatives of the equivalent classes of regular p-adic quadratic forms is obtained as follows.

(a) $n \equiv 1$

Then $q^a(x) = ax^2, a \in M_p$; and the Clifford algebras $C(\mathbf{Q}_p, q^a)$ are isomorphic respectively to $\mathbf{Q}_p \oplus \mathbf{Q}_p, \mathbf{Q}_p[\sqrt{p}], \mathbf{Q}_p[\sqrt{\omega}]$ and $\mathbf{Q}_p[\sqrt{\omega p}]$.

(b) $n \equiv 2$

Then we have over \mathbf{Q}_p^2 (with $\omega = -1$ if $p \equiv 3 \pmod{4}$)

$$\begin{array}{ll} q_0(x) = x_1^2 + x_2^2 & q_4(x) = p x_1^2 + \omega p x_2^2 \text{ if } p \equiv 1 \pmod{4} \\ q_1(x) = x_1^2 + p x_2^2 & \text{(resp. } q_4(x) = p x_1^2 + p x_2^2 \text{ if } p \equiv 3 \pmod{4}) \\ q_2(x) = \omega x_1^2 + \omega p x_2^2 & q_5(x) = x_1^2 + \omega p x_2^2 \\ q_3(x) = x_1^2 + \omega x_2^2 & q_6(x) = p x_1^2 + \omega x_2^2 \end{array}$$

Furthermore $\epsilon(q_\ell) = 1$ if $\ell = 0, 1, 3, 5$ and $\epsilon(q_\ell) = -1$ if $\ell = 2, 4, 6$.

N.B : If $p = 2$, then for $n = 2$, one has

8 regular quadratic forms q such that $\epsilon(q) = 1$
and 7 regular quadratic forms q such that $\epsilon(q) = -1$.

(c) $n \equiv 3$

If (e_1, e_2, e_3) is the canonical basis of \mathbf{Q}_p^3 , then

- $q'_\ell(x) = q_\ell(x_1 e_1 + x_2 e_2) + x_3^2, 0 \leq \ell \leq 6$
and
 - $q'_7(x) = p x_1^2 + \omega x_2^2 + \omega p x_3^2 = q_6(x_1 e_1 + x_2 e_2) + \omega p x_3^2$ if $p \equiv 1 \pmod{4}$
resp.
 - $q'_7(x) = p x_1^2 - x_2^2 + p x_3^2 = q_6(x_1 e_1 + x_2 e_2) + p x_3^2$ if $p \equiv 3 \pmod{4}$
- Furthermore $d(q'_\ell) = d(q_\ell), \epsilon(q'_\ell) = \epsilon(q_\ell), 0 \leq \ell \leq 6$ and $d(q'_7) = -1, \epsilon(q'_7) = -1$.

(d) $n \geq 4$

Let $(e_j)_{1 \leq j \leq n}$ be the canonical basis of \mathbf{Q}_p^n , then

- $q''_\ell(x) = q_\ell(x_1 e_1 + x_2 e_2) + \sum_{j=3}^n x_j^2, 0 \leq \ell \leq 6$.

In other words $q''_\ell(x) = q_\ell(x_1 e_1 + x_2 e_2) + q_0\left(\sum_{j=3}^n x_j e_j\right)$

i.e. $(\mathbf{Q}_p^n, q''_\ell) \simeq (\mathbf{Q}_p^2, q_\ell) \perp (\mathbf{Q}_p^{n-2}, q_0), 0 \leq \ell \leq 6$

and

$$\bullet \quad q''_7(x) = q'_7 \left(\sum_{j=1}^3 x_j e_j \right) + \sum_{j=4}^n x_j^2 = q'_7 \left(\sum_{j=1}^3 x_j e_j \right) + q_0 \left(\sum_{j=4}^n x_j e_j \right)$$

i.e. $(\mathbf{Q}_p^n, q'') \simeq (\mathbf{Q}_p^3, q'_7) \perp (\mathbf{Q}_p^{n-3}, q_0)$

N.B : $p = 2$

If $n = 3$, then the classes of regular quadratic forms have 15 representative forms q' with $\epsilon(q') = 1, \text{resp.} \epsilon(q') = -1$ and $d(q') \neq -1$, obtained from corresponding representative quadratic forms of ranks 2 by adding the rank 1 form x_3^2 . The other representative form is $q'_{15}(x) = -x_1^2 - x_2^2 - x_3^2$ with $\epsilon(q'_{15}) = -1$ and $d(q'_{15}) = -1$.

And if $n \geq 4$, one proceeds as above.

III - 2 The p-adic Clifford algebras

With the above notations , we have the following concrete propositions

Proposition 3 : $p \neq 2$

- (i) $C(\mathbf{Q}_p^2, q_\ell) \simeq M(2, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$.
- (ii) $C(\mathbf{Q}_p^2, q_\ell) \simeq \left(\frac{p, \omega}{\mathbf{Q}_p} \right) = \mathbf{H}_p =$ the p-adic quaternion field , if $\ell = 2, 4, 6$.

Proof

(i) Indeed, if $\ell = 0, 1, 3, 5$; then $\epsilon(q_\ell) = 1$. Therefore $C(\mathbf{Q}_p^2, q_\ell) \simeq M(2, \mathbf{Q}_p)$.

(ii) If $\ell = 2, 4, 6$ then the Clifford algebras $C(\mathbf{Q}_p^2, q_\ell)$ are isomorphic to the quaternion algebras with norm respectively ,

$$N_2(z) = x_0^2 - \omega x_1^2 - \omega p x_2^2 + \omega^2 p x_3^2 ;$$

$$N_4(z) = x_0^2 - p x_1^2 - \omega p x_2^2 + \omega p^2 x_3^2 \text{ if } p \equiv 1 \pmod{4} ;$$

$$\text{(resp. } N_4(z) = x_0^2 - p x_1^2 - p x_2^2 + p^2 x_3^2 \text{ if } p \equiv 3 \pmod{4} \text{)}$$

$$\text{and } N_6(z) = x_0^2 - p x_1^2 - \omega x_2^2 + \omega p x_3^2.$$

It is easily seen that these quadratic forms are anisotropic and equivalent. Therefore $C(\mathbf{Q}_p^2, q_2) \simeq C(\mathbf{Q}_p^2, q_4) \simeq C(\mathbf{Q}_p^2, q_6) \simeq \left(\frac{p, \omega}{\mathbf{Q}_p} \right) = \mathbf{H}_p$ is a skew field. Hence \mathbf{H}_p is the unique quaternion field over \mathbf{Q}_p (according isomorphism). This result obtained directly here is a general result for local fields (cf. [3]).

Proposition 4 : $p \equiv 1 \pmod{4}$

The Clifford algebra $C(\mathbf{Q}_p^3, q'_l)$ is isomorphic to

- (i) $M(2, \mathbf{Q}_p) \oplus M(2, \mathbf{Q}_p)$ if $l = 0$
- (ii) $M(2, \mathbf{Q}_p[\sqrt{p}])$ if $l = 1, 2$
- (iii) $M(2, \mathbf{Q}_p[\sqrt{\omega}])$ if $l = 3, 4$
- (iv) $M(2, \mathbf{Q}_p[\sqrt{\omega p}])$ if $l = 5, 6$
- (v) $\mathbf{H}_p \oplus \mathbf{H}_p$ if $l = 7$

Similarly we have

Proposition 4' : $p \equiv 3 \pmod{4}$

The Clifford algebra $C(\mathbf{Q}_p^3, q'_l)$ is isomorphic to

- (i) $M(2, \mathbf{Q}_p[i])$ if $l = 0, 4$
- (ii) $M(2, \mathbf{Q}_p[\sqrt{-p}])$ if $l = 1, 2$
- (iii) $M(2, \mathbf{Q}_p) \oplus M(2, \mathbf{Q}_p)$ if $l = 3$
- (iv) $M(2, \mathbf{Q}_p[\sqrt{p}])$ if $l = 5, 6$
- (v) $\mathbf{H}_p \oplus \mathbf{H}_p$ if $l = 7$

Proof of Propositions 4 and 4'

Let us recall that if (E, q) is a regular quadratic space over a field K with $n = \dim E$ odd, then $C(E, q) \simeq Z \otimes C_+(E, q)$, where Z is the centre of $C(E, q)$ and $C_+(E, q)$ the subalgebra of even elements. Furthermore, if (e_1, \dots, e_n) is an orthogonal basis of (E, q) then $u = e_1 \dots e_n$ is such that $u^2 = (-1)^{\lfloor \frac{n}{2} \rfloor} d(q)$ and $Z = K[u]$.

In particular for $n = 3$ and $q(x) = \alpha x_1^2 + \beta x_2^2 + \gamma x_3^2$, we have $e_1^2 = \alpha, e_2^2 = \beta, e_3^2 = \gamma; u^2 = -\alpha\beta\gamma = \delta \neq 0$ and $C_+(E, q) = \langle 1, e_1e_2, e_1e_3, e_2e_3 \rangle =$ subspace generated by $1, \dots, e_2e_3$. Put $E_1 = e_1e_2, E_2 = e_1e_3, E_3 = -\alpha e_2e_3$, hence $C_+(E, q) = \langle 1, E_1, E_2, E_3 \rangle$ with $E_1^2 = -\alpha\beta, E_2^2 = -\alpha\gamma, E_1E_2 = E_3 = -E_2E_1$. Therefore $C_+(E, q) \simeq \left(\frac{-\alpha\beta, -\alpha\gamma}{K} \right)$.

Consequently (1) if $\delta \in K^{*2}$, then $Z \simeq K \oplus K$ and $C(E, q) \simeq \left(\frac{-\alpha\beta, -\alpha\gamma}{K} \right) \oplus \left(\frac{-\alpha\beta, -\alpha\gamma}{K} \right)$

(2) if $\delta \notin K^{*2}$, then $Z = K[u]$ is a field and $C(E, q) \simeq \left(\frac{-\alpha\beta, -\alpha\gamma}{K[u]} \right)$.

Applying these remarks to Propositions 4 and 4', one finds the desired isomorphisms. For example if $p \equiv 1 \pmod{4}$ and $l = 2$, then $\delta = -\omega^2 p = (i\omega)^2 p$ and $Z = \mathbf{Q}_p[\sqrt{p}]$, hence

$C(\mathbf{Q}_p^3, q'_2) \simeq \left(\frac{-\omega^2 p, -\omega}{\mathbf{Q}_p[\sqrt{p}]} \right) = \left(\frac{p, \omega}{\mathbf{Q}_p[\sqrt{p}]} \right) \simeq M(2, \mathbf{Q}_p[\sqrt{p}]) : \tilde{q}(v) = p x^2 + \omega y^2$ represents 1 over $\mathbf{Q}_p[\sqrt{p}]$. Also if $\ell = 7$, then $\delta = -p^2 \omega^2 = (i\omega p)^2$, hence $Z \simeq \mathbf{Q}_p \oplus \mathbf{Q}_p$ and since $\left(\frac{-p\omega, -\omega p^2}{\mathbf{Q}_p} \right) \simeq \left(\frac{p\omega, \omega}{\mathbf{Q}_p} \right) \simeq \mathbf{H}_p$ we have $C(\mathbf{Q}_p^3, q'_7) \simeq \mathbf{H}_p \oplus \mathbf{H}_p$.

In the case $p \equiv 3 \pmod{4}$, for example if $\ell = 0$ (resp. $\ell = 3$) we have $\delta = -1$ (resp. $= 1$) and $Z \simeq \mathbf{Q}_p[i]$, (resp. $Z \simeq \mathbf{Q}_p \oplus \mathbf{Q}_p$). Hence $C(\mathbf{Q}_p^3, q'_0) \simeq \left(\frac{-1, -1}{\mathbf{Q}_p[i]} \right) \simeq M(2, \mathbf{Q}_p[i])$, (resp. $C(\mathbf{Q}_p^3, q'_3) \simeq \left(\frac{-1, -1}{\mathbf{Q}_p} \right) \oplus \left(\frac{-1, 1}{\mathbf{Q}_p} \right) \simeq M(2, \mathbf{Q}_p) \oplus M(2, \mathbf{Q}_p)$).

The other verifications are left to the reader.

Lemma 2 : $p \neq 2$

$$C(\mathbf{Q}_p^4, q''_7) \simeq M(2, \mathbf{H}_p). \quad \square$$

Indeed, since $q''_7 = p x_1^2 + \omega x_2^2 + \omega' p x_3^2 + x_4^2$ where $\omega' = \omega$ if $p \equiv 1 \pmod{4}$ and $\omega = -1, \omega' = 1$ if $p \equiv 3 \pmod{4}$; we have $C(\mathbf{Q}_p^4, q''_7) \simeq \left(\frac{p, \omega}{\mathbf{Q}_p} \right) \otimes_2 \left(\frac{\omega' p, 1}{\mathbf{Q}_p} \right) \simeq \mathbf{H}_p \otimes_2 M(2, \mathbf{Q}_p) \simeq M(2, \mathbf{H}_p)$.

Theorem 4 : $p \equiv 1 \pmod{4} ; n \geq 4$

1°) If $n = 2m$, then the Clifford algebra $C(\mathbf{Q}_p^n, q''_\ell)$ is isomorphic to

- (i) $M(2^m, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$
- (ii) $M(2^{m-1}, \mathbf{H}_p)$ if $\ell = 2, 4, 6, 7$

2°) If $n = 2m + 1$, then the Clifford algebra $C(\mathbf{Q}_p^n, q''_\ell)$ is isomorphic to

- (i) $M(2^m, \mathbf{Q}_p) \oplus M(2^m, \mathbf{Q}_p)$ if $\ell = 0$
- (ii) $M(2^m, \mathbf{Q}_p[\sqrt{\tau}])$ if $\ell = 1, 2, 3, 4, 5, 6$

with $\tau = p$ (resp. $\omega, \text{resp. } \omega p$) for $\ell = 1, 2$ (resp. $\ell = 3, 4$; resp. $5, 6$).

- (iii) $M(2^{m-1}, \mathbf{H}_p) \oplus M(2^{m-1}, \mathbf{H}_p)$ if $\ell = 7$

Proof :

1°) $n = 2m$

Notice that $C(\mathbf{Q}_p^n, q''_\ell) \simeq C(\mathbf{Q}_p^{n-2}, q_0) \otimes_2 C(\mathbf{Q}_p^2, q_\ell), 0 \leq \ell \leq 6$. But by Proposition 3, we have $C(\mathbf{Q}_p^2, q_\ell) \simeq M(2, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^2, q_\ell) \simeq \mathbf{H}_p$ if $\ell =$

2, 4, 6. Since $C(\mathbf{Q}_p^{n-2}, q_0) \simeq M(2^{m-1}, \mathbf{Q}_p)$ by Theorem 1 - (i) - , we have $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{m-1}, \mathbf{Q}_p) \otimes_2 M(2, \mathbf{Q}_p) \simeq M(2^m, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{m-1}, \mathbf{Q}_p) \otimes_2 \mathbf{H}_p \simeq M(2^{m-1}, \mathbf{H}_p)$ if $\ell = 2, 4, 6$.

For $\ell = 7$, applying Lemma 2 and Theorem 1 - (i) - we obtain $C(\mathbf{Q}_p^n, q_\ell'') \simeq C(\mathbf{Q}_p^{n-4}, q_0) \otimes_2 C(\mathbf{Q}_p^4, q_\ell'') \simeq M(2^{m-2}, \mathbf{Q}_p) \otimes_2 M(2, \mathbf{H}_p) \simeq M(2^{m-1}, \mathbf{H}_p)$.

2°) $n = 2m+1$

If $1 \leq \ell \leq 6$, then we have $C(\mathbf{Q}_p^n, q_\ell'') \simeq C(\mathbf{Q}_p^{n-3}, q_0) \otimes_2 C(\mathbf{Q}_p^3, q_\ell') \simeq M(2^{m-1}, \mathbf{Q}_p) \otimes_2 C(\mathbf{Q}_p^3, q_\ell')$.

Applying Proposition 4, we obtain the isomorphism $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^m, \mathbf{Q}_p[\sqrt{\tau}])$ as claimed.

The case $\ell = 0$ is Theorem 1 - (ii) -

If $\ell = 7$, then $C(\mathbf{Q}_p^3, q_7') \simeq \mathbf{H}_p \oplus \mathbf{H}_p$ and $C(\mathbf{Q}_p^n, q_7'') \simeq M(2^{m-1}, \mathbf{Q}_p) \otimes_2 (\mathbf{H}_p \oplus \mathbf{H}_p) \simeq M(2^{m-1}, \mathbf{H}_p) \oplus M(2^{m-1}, \mathbf{H}_p)$.

Theorem 5 : $p \equiv 3(mod.4) ; n \geq 4$

The Clifford algebra $C(\mathbf{Q}_p^n, q_\ell'')$ is isomorphic to the following matrix algebra or direct sum of two matrix algebras.

1°) $n = 4m$

- (i) $M(2^{2m}, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$
- (ii) $M(2^{2m-1}, \mathbf{H}_p)$ if $\ell = 2, 4, 6, 7$

2°) $n = 4m+1$

- (i) $M(2^{2m}, \mathbf{Q}_p) \oplus M(2^{2m}, \mathbf{Q}_p)$ if $\ell = 0, 4$
- (ii) $M(2^{2m}, \mathbf{Q}_p[\sqrt{\tau}])$ if $\ell = 1, 2, 3, 5, 6, 7$

with $\tau = p$ (resp. $-1, res. -p$) for $\ell = 1, 2$ (resp. $\ell = 3, 7, resp. \ell = 5, 6$).

3°) $n = 4m+2$

- (i) $M(2^{2m+1}, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$
- (ii) $M(2^{2m}, \mathbf{H}_p)$ if $\ell = 2, 4, 6, 7$

$$4^{\circ) \quad \underline{n = 4m+3}$$

$$(i) \quad M(2^{2m+1}, \mathbf{Q}_p) \oplus M(2^{2m+1}, \mathbf{Q}_p) \quad \text{if} \quad \ell = 3$$

$$(ii) \quad M(2^{2m+1}, \mathbf{Q}_p[\sqrt{\tau}]) \quad \text{if} \quad \ell = 0, 1, 2, 4, 5, 6,$$

with $\tau = -1$ (resp. $-p, \text{res. } p$) for $\ell = 0, 4$ (resp. $\ell = 1, 2, \text{resp. } \ell = 5, 6$).

$$(iii) \quad M(2^{2m}, \mathbf{H}_p) \oplus M(2^{2m}, \mathbf{H}_p) \quad \text{if} \quad \ell = 7.$$

Proof :

$$1^{\circ) \quad \underline{n = 4m}$$

As in Lemma 2, it is readily seen that $C(\mathbf{Q}_p^4, q_\ell'') \simeq M(2^2, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^4, q_\ell'') \simeq M(2, \mathbf{H}_p)$ if $\ell = 2, 4, 6, 7$.

If $n = 4m$, $m \geq 2$, we have $C(\mathbf{Q}_p^n, q_\ell'') \simeq C(\mathbf{Q}_p^{n-4}, q_0) \otimes_2 C(\mathbf{Q}_p^4, q_\ell'')$. But Theorem 3 - (i) - gives $C(\mathbf{Q}_p^{n-4}, q_0) \simeq M(2^{2m-2}, \mathbf{Q}_p)$. Therefore $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{2m-2}, \mathbf{Q}_p) \otimes_2 M(2^2, \mathbf{Q}_p) \simeq M(2^{2m}, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{2m-2}, \mathbf{Q}_p) \otimes_2 M(2^2, \mathbf{H}_p) \simeq M(2^{2m-1}, \mathbf{H}_p)$ if $\ell = 2, 4, 6, 7$.

$$2^{\circ) \quad \underline{n = 4m+1}$$

With notations used in the proof of Propositions 4 and 4' we have $C(\mathbf{Q}_p^n, q_\ell'') \simeq Z \otimes C_+(\mathbf{Q}_p^n, q_\ell'')$ and $Z = \mathbf{Q}_p[u]$ where $u^2 = d(q_\ell'')$. Hence Z is isomorphic to $\mathbf{Q}_p \oplus \mathbf{Q}_p$ if $\ell = 0, 4$; resp. $\mathbf{Q}_p[\sqrt{p}]$ if $\ell = 1, 2$; resp. $\mathbf{Q}_p[\sqrt{-1}]$ if $\ell = 3, 7$; resp. $\mathbf{Q}_p[\sqrt{-p}]$ if $\ell = 5, 6$. On the other hand $C_+(\mathbf{Q}_p^n, q_\ell'') \simeq C_+(\mathbf{Q}_p \cdot x_n, x_n^2) \otimes C(\mathbf{Q}_p^{n-1}, -q_\ell'') \simeq C(\mathbf{Q}_p^{n-1}, -q_\ell'') \simeq M(2^{2m}, \mathbf{Q}_p)$. Hence $C(\mathbf{Q}_p^n, q_\ell'') \simeq Z \otimes M(2^{2m}, \mathbf{Q}_p)$ which proves the isomorphisms.

$$3^{\circ) \quad \underline{n = 4m+2}$$

Since $n - 2 = 4m$, we obtain $C(\mathbf{Q}_p^n, q_\ell'') \simeq C(\mathbf{Q}_p^{4m}, q_0) \otimes_2 C(\mathbf{Q}_p^2, q_\ell)$.

By Theorem 2 - (i) - one has $C(\mathbf{Q}_p^{4m}, q_0) \simeq M(2^{2m}, \mathbf{Q}_p)$ and by Proposition 3, $C(\mathbf{Q}_p^2, q_\ell) \simeq M(2, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^2, q_\ell) \simeq \mathbf{H}_p$ if $\ell = 2, 4, 6$. It follows that $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{2m+1}, \mathbf{Q}_p)$ if $\ell = 0, 1, 3, 5$ and $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{2m}, \mathbf{H}_p)$ if $\ell = 2, 4, 6$.

For the case $\ell = 7$, since $n - 4 = 4(m - 1) + 2$ we have $C(\mathbf{Q}_p^n, q_\ell'') \simeq C(\mathbf{Q}_p^{n-4}, q_0) \otimes_2 C(\mathbf{Q}_p^4, q_\ell'')$. By theorem 2 - (iii) -, $C(\mathbf{Q}_p^{n-4}, q_0) \simeq M(2^{2m+1}, \mathbf{Q}_p)$ and by Lemma 2, $C(\mathbf{Q}_p^4, q_\ell'') \simeq M(2, \mathbf{H}_p)$. Hence $C(\mathbf{Q}_p^n, q_\ell'') \simeq M(2^{2m}, \mathbf{H}_p)$.

Notice that in 1°) and 3°) the exponent of 2 is $\frac{n}{2}$.

4°) $\underline{n} = 4m+3$

Here, $n-3 = 4m$ and $C(\mathbf{Q}_p^n, q''_l) \simeq C(\mathbf{Q}_p^{4m}, q_0) \otimes_2 C(\mathbf{Q}_p^3, q'_l)$. But $C(\mathbf{Q}_p^{4m}, q_0) \simeq M(2^{2m}, \mathbf{Q}_p)$ and by Proposition 4', $C(\mathbf{Q}_p^3, q'_l)$ is isomorphic to $M(2, \mathbf{Q}_p) \oplus M(2, \mathbf{Q}_p)$ if $\ell = 3$, resp. $\mathbf{H}_p \oplus \mathbf{H}_p$ if $\ell = 7$, resp. $M(2, \mathbf{Q}_p[\sqrt{\tau}])$ if $\ell = 0, 1, 2, 4, 5, 6$ with $\tau = -1$ for $\ell = 0, 4$; $\tau = -p$ for $1, 2$ and $\tau = p$ for $\ell = 5, 6$.

Taking tensor product we obtain the desired isomorphisms.

Remark :

As for $C(\mathbf{Q}_p^n, q_0)$, for the other Clifford algebras $C(\mathbf{Q}_p^n, q''_l)$ we have 2-periodicity when $p \equiv 1 \pmod{4}$ and 4-periodicity when $p \equiv 3 \pmod{4}$.

N.B. When $p = 2$, in the same way one can give as above the table of the 2-adic Clifford algebras.

REFERENCES

- [1] R. DEHEUVELS, *Formes quadratiques et groupes classiques*, P.U.F. - Paris - 1981.
- [2] B. DIARRA, *Remarques sur les matrices orthogonales (resp. symétriques) à coefficients p-adiques*, Ann. Sci. "Blaise Pascal", Clermont II, sér. Math., Fasc. 26è - 1990 - p. 31-50.
- [3] T.Y. LAM, *The algebraic theory of quadratic forms*, W.C. Benjamin, Inc. - Reading, Massachusetts - 1973.
- [4] P. RIBENBOIM, *L'arithmétique des corps*, Hermann - Paris - 1972.
- [5] J.P. SERRE, *Cours d'arithmétique*, P.U.F. - Paris - 1970.